

Cyclophosphamide

Methylprednisolone

. . . * . .

= Abstract =

Effect of Cyclophosphamide and High Dose Methylprednisolone on Paraquat Poisoned Rats

Uk Jin Kim, M.D., Sung Pil Chung, M.D., Yoo Shuk Park, M.D.,
Min Goo Lee, M.D.*, Seung Ho Kim, M.D., Hahn Shick Lee, M.D.

Department of Emergency Medicine, and Pharmacology*
Yonsei University College of Medicine

Background: Paraquat(PQ) is used widely all over the world for its excellent effect as a herbicide. But its mortality rate is known to be very high, because there is no effective therapeutic modality. Recently, surprising improvement in survival rate was reported using the cyclophosphamide & methylprednisolone pulse therapy in paraquat poisoning. However, this report was not based upon animal study, we designed this experiment to confirm the therapeutic effect.

Methods: Under the halothane anesthesia, paraquat dichloride 40 mg/kg was injected intraperitoneally to 18 Sprague-Dawley rats. Two hours later, cyclophosphamide 40 mg/kg IP and methylprednisolone 62.5 mg/kg IM were injected in the treatment group(n=9). After 24 hours, we examined serum creatinine levels and pathologic findings of lung stained with hematoxylin-eosin and Masson's trichrome. And 72 hour mortality was compared between 2 groups(5 rats respectively).

Results: There were no statistical differences between the treatment group and control group in serum creatinine level, degree of lung injury, and survival rates.

Conclusion: Cyclophosphamide and high dose methylprednisolone combination therapy did not decrease pulmonary toxicity and mortality of paraquat poisoned rats. Further animal studies using various doses and administrative methods of above medications are necessary to demonstrate their effects.

Key Words: Paraquat poisoning, Cyclophosphamide, Methylprednisolone

I. cyclophosphamide methyl-
prednisolone ,
(1,1'-dimethyl-4,4' dipyridylum) 40% 가
19 , 1955 23).
가
가
1300
1,2) 1970 Lin cyclophosphamide
가 1000 methylprednisolone
3).
가
가 NADPH II.
superoxide (O₂), hydroperoxy (HO₂)
(H₂O₂) 470-540 g (Sprague-Dawley)
18 9
cyclophosphamide methylpre-
dnisolone 1
20% 15 ml 2 .
100% 5). 가
2
6-10) 11) , 1 2 ,
가 가
12).
5%
, Bentonite 7,13,14)
70%
, 1 2
13,15,16) 가 50%
17) superoxide dismutase 18,19)가 (Zeneca agrochemicals, Fernfurst, UK)
. prednisone azathioprine 50 40 mg/kg(
가 가 29.9 mg/kg) . 2
20). cyclophosphamide dexametha-
sone 75% . 2
21) cyclophosphamide 40 mg/kg
22). methylprednisolone 62.5 mg/kg
Lin 2
,

24

6 ml

(

1, 2).

10%

7


Hematoxylin and eosin Masson and
trichrome

(5), (4), (3),
(2), (1) 5

10

10

12



1. : (×
100, H-E)


SPSS

8.0

III.

2 가 4.3
mg/dl, 2.8 mg/dl 가 , 18

(1).




2. : (×100, H-E)


(1).

1.	cyclophosphamide	methylprednisolone	p *
(mg/dl)	1.17 ± 1.40	0.58 ± 0.38	>0.05
**	2.89 ± 0.60	3.33 ± 0.70	>0.05
**	1.22 ± 0.44	1.44 ± 0.73	>0.05
**	2.67 ± 0.50	2.33 ± 0.50	>0.05
**	2.88 ± 0.93	2.56 ± 0.53	>0.05

*, **, , **.




3. : (×400, H-E)



5. Cyclophosphamide methylprednisolone

($P>0.05$) (: , :) .



4. : (×400, Masson's trichrome)

가
superoxide , peroxide

가

12)

가 fibronectin

II 가

가

9)

IV.

. Dimethyl sulfoxide hydroxyl

(1,1'-dimethyl-4,4'-dipyridylum)

²⁴⁾ Xantine oxidase inhibitor

1960 , 100

allopurinol

	25) . Superoxide dismutase	prednisolone	가 .
(SOD)가	18)	Lin	가
19) .	cyclophosphamide	가 .	cyclophosphamide
D-penicillamine		methylprednisolone	,
가	26) .		
steroid			.
20) cyclophosphamide	dexame-		,
thasone	75%		가 가 가
21)	가		
22) .		가 ,	
		가	.
cyclophosphamide methylprednisolone		18.2%	
	40%	가	,
가	23) .		.
,	,		
		V.	
		cyclophosphamide	
		methylprednisolone	
cyclophosphamide methylprednisolone		1. cyclophosphamide	methylpre-
		dnisolone	
	2		.
		2. cyclophosphamide	methylpre-
		dnisolone	
가			.
가	가	cyclophosphamide	
가		methylprednisolone	가
가			.
가			
cyclophosphamide	methyl-	1. Crome P: Paraquat poisoning. <i>Lancet</i> 8:333-334, 1986.	
		2. Yamashita M, Matsuo H, Ando Y: Analysis of	

- 1,000 consecutive cases of acute poisoning in the suburb of Tokyo leading to hospitalization. *Vet Human Toxicol* 38:34-35, 1996.
3. : .
3319 : 16 , 1999.
4. Bus JS, Cagen SZ, Olgaard M, Gibson JE: A mechanism of paraquat toxicity in mice and rats. *Toxicol Appl Pharmacol* 35:501-513, 1976.
5. Ellenhorn MJ: paraquat. 2nd Ed. Ellenhorn 's Medical Toxicology: Diagnosis and treatment of human poisoning. *Williams & Wilkins*, p1631-1637, 1997.
6. Kimbrough RD, Gaines TB: Toxicity of paraquat to rats and its effect on rat lungs. *Toxicol Appl Pharmacol* 17:679-697, 1970.
7. Smith P, Heath D, Kay JM: The pathogenesis and structure of paraquat-induced pulmonary fibrosis in rats. *J pathol* 114:57-67, 1973.
8. Greenberg DB, Reiser KM, Last JA: Correlation of biochemical and morphologic manifestations of acute pulmonary fibrosis in rats administered paraquat. *Chest* 74:421-425, 1978.
9. Fukuda Y, Ferrans VJ, Schoenberger CI, Rennard SI, Crystal RG: Patterns of pulmonary structural remodeling after experimental paraquat toxicity. The morphogenesis of intraalveolar fibrosis. *Am J pathol* 118:452-475, 1985.
10. Schoenberger CI, Rennard SI, Bitterman PB, Fukuda Y, Fermans VJ, Crystal RG: Paraquat-induced pulmonary fibrosis; role of the alveolitis in modulating the development of fibrosis. *Am Rev Respir Dis* 129:168-173, 1984.
11. Ecker JL, Hook JB, Gibson JE: Nephrotoxicity of paraquat in mice. *Toxicol Appl Pharmacol* 34:178-186, 1975.
12. Copland GM, Kolin A, Shulman HS: Fatal pulmonary intra-alveolar fibrosis after paraquat ingestion. *N Engl J Med* 291:290-292, 1974.
13. Fairshter RD, Wilson AF: Paraquat poisoning; Manifestations and therapy. *Am J Med* 59:751-753, 1975.
14. Idid SZ, Lee CY: Effects of Fuller 's Earth and activated charcoal on oral absorption of paraquat in rabbits. *Clin Experi Pharmacol & Physiol* 23:679-681, 1996.
15. Solfrank G, Mathes C, Clarmann V: Hemoperfusion through activated charcoal in paraquat intoxication. *Acta Pharmacol Toxicol(Suppl)* 41:91-101, 1977.
16. Vale JA, Crome P, Volans GN, Widdop B, Goulding R: The treatment of paraquat poisoning using oral sorbents and charcoal hemoperfusion. *Acta Pharmacol Toxicol(suppl)* 41:109-117, 1977.
17. Rhodes ML, Zavala DC, Brown D: Hypoxic protection in paraquat poisoning. *Lab invest* 35: 496-500, 1976.
18. Wasserman B, Block ER: Prevention of acute paraquat toxicity in rats by superoxide dismutase. *Aviat Space Environ Med* 49:805-809, 1978.
19. Patterson CE, Rhodes ML: The effect of superoxide dismutase on paraquat mortality in mice and rats. *Toxicol appl pharmacol* 62:65-72, 1982.
20. Malcolmson E, Beesley J: Unsuccessful immuno-suppressant treatment of paraquat poisoning(letter). *Brit Med J* 650-651, 1975.
21. Addo E, Ramandial S, Poon-King T: High dosage cyclophosphamide and dexamethasone treatment of paraquat poisoning with 75% survival. *West Indian Med J* 33:220-226, 1984.
22. Perriens JH, Benimadho S, Kiauw LI, Wisse J, Chee H: High-dose cyclophosphamide and dexamethasone in paraquat poisoning: a prospective study. *Hum Exp Toxicol* 11:129-134, 1992.
23. Lin JL, Leu ML, Liu YC, Chen GH: Sprospective clinical trial of pulse therapy with glucocorticoid and cyclophosphamide in moderate to severe paraquat-poisoned patients. *Am J Respir Crit care Med* 159:357-360, 1999.
24. , , , :
가
36(1): 51-64, 1995.
25. , , , , :
.
20(1):1-12, 1995.
26. , : 가
.
7:458-471, 1996.